

**THE CHINESE UNIVERSITY OF HONG KONG**  
**Department of Mathematics**  
**2018-2019 semester 1 MATH4060**  
**week 5 tutorial**

Underlined contents were not included in the tutorial because of time constraint, but included here for completeness.

Below is a brief introduction to properties of harmonic functions. Removable singularity theorem and Liouville's theorem for harmonic functions are proven by maximum principle and Poisson integral formula. The main reference is Chapter 2 of Gilbarg and Trudinger's *Elliptic Partial Differential Equations of Second Order*. Below,  $\Omega$  always denotes a nonempty connected open set in  $\mathbb{R}^2 = \mathbb{C}$ .

## 1 Properties of Harmonic Functions

A  $C^2$  function  $u : \Omega \rightarrow \mathbb{R}$  is harmonic iff  $\Delta u = u_{xx} + u_{yy} = 0$ .

Harmonic functions and holomorphic functions are intimately related.

1.  $f$  is holomorphic iff  $\partial_{\bar{z}}f = 0$ , whereas  $u$  is harmonic iff  $\partial_z\partial_{\bar{z}}u = 0$ .
2. If  $f$  is holomorphic, then  $\Re f$ ,  $\Im f$  and  $\log |f|$  are harmonic whenever finitely defined. If  $\Omega$  is simply connected and  $u$  is harmonic, then  $f = u + iv$ , where  $v = \int(u_x dy - u_y dx)$ , is holomorphic, and  $\log |e^f| = u$ .
3. (Cauchy integral formul and mean-value property) If  $f$  is holomorphic, then

$$f(z) = \frac{1}{2\pi i} \int_{\partial B(z,r)} \frac{f(w)}{w-z} dw.$$

If  $u$  is harmonic, then

$$u(z) = \frac{1}{2\pi r} \int_{\partial B(z,r)} u(w) dw = \frac{1}{2\pi} \int_{\partial B(z,r)} \frac{u(w)}{|w-z|} dw. \quad (1)$$

4. (strong maximum (modulus) principle) If a holomorphic  $f$  attains the maximum modulus in the interior, then it is constant. If a harmonic  $u$  attains the maximum in the interior, then it is constant.
5. (weak maximum (modulus) principle) The maximum modulus of a holomorphic function or a harmonic function on a bounded domain is attained on the boundary.

Mean-value property for harmonic function is more rigid than that for holomorphic function because the domain of integration in (1) cannot be any  $\partial B(w,r)$  containing  $z$ . Indeed, the offset mean-value property is given by the more involved Poisson integral formula.

**Proposition 1** (Poisson integral formula). Suppose  $u$  is harmonic on a neighbourhood of  $\overline{B(0, R)}$ . Let  $\varphi = u|_{\partial B(0, R)}$ . Then for  $x \in B(0, R)$

$$u(x) = \int_{\partial B(0, R)} \varphi(y) P_2(x, y) dy, \quad (2)$$

where  $P_n(x, y) = \frac{1}{|\partial B(0, R)|} \frac{R^2 - |x|^2}{R^2} \left( \frac{R}{|x-y|} \right)^n$ .

Conversely, if  $\varphi$  is a continuous function on  $\partial B(0, R)$ , then (2) defines a harmonic function on  $B(0, R)$  whose continuous extension to  $\partial B(0, R)$  exists and agrees with  $\varphi$ .

**Corollary 2.** Harmonic functions are smooth.

Below, we prove removable singularity theorem and Liouville's theorem for harmonic functions.

**Proposition 3** (Removable singularity theorem). Suppose  $u$  is harmonic on  $B(0, r) \setminus \{0\}$ . If  $u(z) = o(\log |z|)$  as  $z \rightarrow 0$ , then  $u$  extends to a harmonic function on  $B(0, r)$ .

*Proof.* It suffices to show  $u$  agrees to  $\tilde{u}$  defined by Poisson integral formula, which is a harmonic function on  $B(0, r)$ . Let  $w = \tilde{u} - u$ . Then  $w(z) = o(\log |z|) = o(\log |z| - \log r)$ . Note that both  $w$  and  $\log |z| - \log r$  vanish on  $\partial B(0, r)$ . By maximum principle, for  $\varepsilon > 0$ , since  $\pm w(z) + \varepsilon \log |z| \rightarrow -\infty$ ,  $\sup_{B(0, r) \setminus \{0\}} \pm w + \varepsilon(\log |z| - \log r) \leq 0$ . The result follows by letting  $\varepsilon \rightarrow 0$ .  $\square$

To prove Liouville's property, it is handy to have an estimate on the gradient.

**Proposition 4** (gradient estimate). Suppose  $u$  is harmonic on a neighbourhood of  $\overline{B(0, R)}$ . Then

$$|\partial_i u(0)| \leq \frac{n}{R} \|u\|_{L^\infty(B(0, R))}.$$

*Remark.* Repeated application of the gradient estimate shows harmonic functions are in fact analytic.

*Proof.* Apply differentiation under the integral sign on Poisson integral formula.  $\square$

**Proposition 5** (Liouville's theorem). If a harmonic function on  $\mathbb{R}^2$  is bounded, then it is constant.

*Proof.* Let  $R \rightarrow \infty$  in the gradient estimate.  $\square$

**Exercise 6.** Complete the following alternative proof of Liouville's theorem:

By Poisson integral formula, we have the following Harnack inequality for nonnegative harmonic  $u$  on  $\mathbb{R}^2$

$$\frac{1}{(R + |x|)} \frac{R - |x|}{R} u(0) \leq u(x) \leq \frac{1}{(R - |x|)} \frac{R + |x|}{R} u(0).$$

Liouville's theorem for nonnegative functions then follows by letting  $R \rightarrow \infty$  on the far right. The general case follows by translation.