

CMSC5724: Exercise List 3

Problem 1. Let P be a set of 4 points: $A = (1, 2, 1)$, $B = (2, 1, 1)$, $C = (0, 1, 1)$ and $D = (1, 0, 1)$. A and B have label 1, while C and D have label -1 . Execute Perceptron on P . Give the weight vector \mathbf{w} maintained by the algorithm after each iteration.

Answer. At the beginning, $\mathbf{w} = (0, 0, 0)$. We use \mathbf{A} to denote the vector form of A . Define \mathbf{B} , \mathbf{C} , and \mathbf{D} similarly.

Iteration 1. Since \mathbf{A} does not satisfy $\mathbf{A} \cdot \mathbf{w} > 0$, update \mathbf{w} to $\mathbf{w} + \mathbf{A} = (0, 0, 0) + (1, 2, 1) = (1, 2, 1)$.

Iteration 2. Since \mathbf{C} does not satisfy $\mathbf{C} \cdot \mathbf{w} < 0$, update \mathbf{w} to $\mathbf{w} - \mathbf{C} = (1, 2, 1) - (0, 1, 1) = (1, 1, 0)$.

Iteration 3. Since \mathbf{C} does not satisfy $\mathbf{C} \cdot \mathbf{w} < 0$, update \mathbf{w} to $\mathbf{w} - \mathbf{C} = (1, 1, 0) - (0, 1, 1) = (1, 0, -1)$.

Iteration 4. Since \mathbf{A} does not satisfy $\mathbf{A} \cdot \mathbf{w} > 0$, update \mathbf{w} to $\mathbf{w} + \mathbf{A} = (1, 0, -1) + (1, 2, 1) = (2, 2, 0)$.

Iteration 5. Since \mathbf{C} does not satisfy $\mathbf{C} \cdot \mathbf{w} < 0$, update \mathbf{w} to $\mathbf{w} - \mathbf{C} = (2, 2, 0) - (0, 1, 1) = (2, 1, -1)$.

Iteration 6. Since \mathbf{C} does not satisfy $\mathbf{C} \cdot \mathbf{w} < 0$, update \mathbf{w} to $\mathbf{w} - \mathbf{C} = (2, 1, -1) - (0, 1, 1) = (2, 0, -2)$.

Iteration 7. Since \mathbf{A} does not satisfy $\mathbf{A} \cdot \mathbf{w} > 0$, update \mathbf{w} to $\mathbf{w} + \mathbf{A} = (2, 0, -2) + (1, 2, 1) = (3, 2, -1)$.

Iteration 8. Since \mathbf{C} does not satisfy $\mathbf{C} \cdot \mathbf{w} < 0$, update \mathbf{w} to $\mathbf{w} - \mathbf{C} = (3, 2, -1) - (0, 1, 1) = (3, 1, -2)$.

Iteration 9. Since \mathbf{D} does not satisfy $\mathbf{D} \cdot \mathbf{w} < 0$, update \mathbf{w} to $\mathbf{w} - \mathbf{D} = (3, 1, -2) - (1, 0, 1) = (2, 1, -3)$.

Iteration 10. No more violation points.


Problem 2. Let P be a set of multidimensional points where each point has a label equal to 1 or -1 . We want to design an algorithm to achieve the following purpose:

- Either return a separation plane (see the lecture notes for the definition of separation plane);
- Or declare that P has no separation planes with a margin at least γ .

Your algorithm must still work even if no separation planes exist.

Answer. Run Perceptron and return whatever plane found by the algorithm. If the algorithm still has not finished after R^2/γ^2 corrections, force it to stop and declare that no separation plane has a margin at least γ .

Problem 3. Consider the set of points below where points of different colors carry different labels. Only two points have their coordinates shown. Apply Perceptron to find a separation plane on the set. Prove: Perceptron finishes after at most 5 iterations.



Answer. The y-axis is a separation plane with margin $\gamma = 1$. Clearly, the largest distance from a point to the origin is $R = \sqrt{5}$. Hence, Perceptron performs at most $R^2/\gamma^2 = 5$ iterations.

Problem 4. Some people prefer the following variant of the Perceptron algorithm:

1. $\mathbf{w} = \mathbf{0}$
2. **while** there is a violating point \mathbf{p}
3. **if** \mathbf{p} has label 1
4. $\mathbf{w} = \mathbf{w} + \lambda \cdot \mathbf{p}$
- else**
5. $\mathbf{w} = \mathbf{w} - \lambda \cdot \mathbf{p}$

where λ is a positive real-value constant. In the version we discussed in the lecture, $\lambda = 1$. Prove: regardless of λ , Perceptron always terminates in R^2/γ^2 iterations, where R is the maximum distance of the points to the origin and γ the largest margin of all separation planes.

Answer. Let \mathbf{w}_k denote the vector \mathbf{w} after the k -th iteration. Following the analysis we discussed in the lecture, we can prove the two inequalities below:

$$\begin{aligned} |\mathbf{w}_k| &\geq \lambda \cdot k \cdot \gamma \\ |\mathbf{w}_k|^2 &\leq k \cdot \lambda^2 \cdot R^2 \end{aligned}$$

The two inequalities give $\lambda^2 \cdot k^2 \cdot \gamma^2 \leq k \cdot \lambda^2 \cdot R^2$, which indicates $k \leq R^2/\gamma^2$.

Problem 5. Let P be a set of points in \mathbb{R}^d , where each point is labeled 1 or -1 . A d -dimensional plane π is a *separation plane* of P if

- π does not pass any point in P ;
- the points of the two labels in P fall on different sides of π .

Note that we do not require π to pass the origin.

Construct a $(d + 1)$ -dimensional point set P' as follows: given each $p \in P$, add to P' the point $(p[1], p[2], \dots, p[d], 1)$ (i.e., adding a new coordinate 1), carrying the same label as p . Prove: P has a separation plane if and only if P' has a separation plane passing the origin of \mathbb{R}^{d+1} .

Answer: Given a point $p \in P$, we use \mathbf{p} to denote its vector form. Similarly, we use \mathbf{p}' to denote the vector form of a point $p' \in P'$.

Only-if direction. Suppose that P has a separation plane. Then, there must be a d -dimensional plane $\mathbf{w} \cdot \mathbf{x} + w_{d+1} = 0$ such that for every $p \in P$:

$$\begin{cases} \mathbf{w} \cdot \mathbf{p} + w_{d+1} > 0 & \text{if } p \text{ has label } 1 \\ \mathbf{w} \cdot \mathbf{p} + w_{d+1} < 0 & \text{if } p \text{ has label } -1. \end{cases} \quad (1)$$

Let $\mathbf{w}' = (\mathbf{w}[1], \mathbf{w}[2], \dots, \mathbf{w}[d], w_{d+1})$. Every $p' \in P'$ has the same label as $(p'[1], p'[2], \dots, p'[d])$ in P and $p'[d+1] = 1$. From (1), we have

- $\mathbf{w}' \cdot \mathbf{p}' = \sum_{i=1}^d \mathbf{w}[i]p'[i] + w_{d+1} > 0$ if p' has label 1;
- $\mathbf{w}' \cdot \mathbf{p}' = \sum_{i=1}^d \mathbf{w}[i]p'[i] + w_{d+1} < 0$ if p' has label -1 .

Hence, P' has a separation plane (i.e., $\mathbf{w}' \cdot \mathbf{x} = 0$) passing the origin of \mathbb{R}^{d+1} .

If-direction. Suppose that P' has a separation plane passing the origin of \mathbb{R}^{d+1} . Then, there must be a $(d+1)$ -dimensional vector \mathbf{w}' such that for every $p' \in P'$:

$$\begin{cases} \mathbf{w}' \cdot \mathbf{p}' > 0 & \text{if } p' \text{ has label } 1 \\ \mathbf{w}' \cdot \mathbf{p}' < 0 & \text{if } p' \text{ has label } -1. \end{cases} \quad (2)$$

Let $\mathbf{w} = (\mathbf{w}'[1], \mathbf{w}'[2], \dots, \mathbf{w}'[d])$. Every $p \in P$ has the same label as $p' = (p[1], p[2], \dots, p[d], 1)$ in P' . From (2), we know

- $\mathbf{w} \cdot \mathbf{p} + \mathbf{w}'[d+1] = \mathbf{w}' \cdot \mathbf{p}' > 0$ if p has label 1;
- $\mathbf{w} \cdot \mathbf{p} + \mathbf{w}'[d+1] = \mathbf{w}' \cdot \mathbf{p}' < 0$ if p has label -1 .

It thus follows that P has a separation plane $\mathbf{w} \cdot \mathbf{x} + \mathbf{w}'[d+1] = 0$.