

Lecture 12: Recall:

Prop: Let T be a linear operator on a vector space V and let λ be an eigenvalue of T . Then, $\vec{v} \in V$ is an eigenvector of T corresponding to λ iff:

$$\vec{v} \in N(T - \lambda I_V) \setminus \{\vec{0}\}$$

Def: Let T be a linear operator on a vector space V and let λ be an eigenvalue of T .

Then: the subspace $E_\lambda := \begin{matrix} \text{def} \\ N(T - \lambda I_V) = \{ \vec{x} \in V : T(\vec{x}) = \lambda \vec{x} \} \end{matrix} \subset V$ is called the eigenspace of T corresponding to λ .

Eigenspace of a matrix $A \in M_{n \times n}(F)$ is defined as those of L_A

Prop: Let T be a linear operator on a vector space V , and let $\lambda_1, \lambda_2, \dots, \lambda_k$ be distinct eigenvalues of T . If $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k$ are eigenvectors of T corresponding to $\lambda_1, \lambda_2, \dots, \lambda_k$ respectively, then $\{\vec{v}_1, \dots, \vec{v}_k\}$ are linearly independent.

Corollary: A linear operator T on an n -dim vector space V which has n distinct eigenvalues is diagonalizable.

Proof: Let $\vec{v}_1, \dots, \vec{v}_n \in V$ be the eigenvectors corresponding to n distinct eigenvalues. Then, the prop. says $\{\vec{v}_1, \dots, \vec{v}_n\}$ is lin. independent. $\therefore \{\vec{v}_1, \dots, \vec{v}_n\}$ forms a basis of eigenvectors. $\therefore T$ is diagonalizable.

Def: Let λ be an eigenvalue of a linear operator or matrix with characteristic polynomial $f(t)$. The **algebraic multiplicity** of λ , denoted $M_T(\lambda)$ or $M_A(\lambda)$ is the multiplicity of λ as a zero of $f(t)$, i.e. the largest positive integer k s.t. $(t-\lambda)^k \mid f(t)$.

Example: • 1 is eigenvalue of $I_V: V \rightarrow V$

with $\mu_{I_V}(1) = \dim(V)$

$$f(t) = \det \left([I_V]_\beta - t I_n \right) = \begin{pmatrix} 1-t & & & \\ & 1-t & & \\ & & \ddots & \\ & & & 1-t \end{pmatrix} = (1-t)^n$$

$$\bullet A = \begin{pmatrix} 3 & 1 & 0 \\ 0 & 3 & 4 \\ 0 & 0 & 5 \end{pmatrix} . \quad f_A(t) = (3-t)^2(5-t)$$

$$\mu_A(3) = 2, \quad \mu_A(5) = 1$$

Prop: Let T be a linear operator on a finite-dim vector space V and let λ be an eigenvalue of T with algebraic multiplicity $M_T(\lambda)$. Then:

$$1 \leq \dim(E_\lambda) \leq M_T(\lambda)$$

We call $\gamma_T(\lambda) \stackrel{\text{def}}{=} \dim(E_\lambda)$ the geometric multiplicity of λ .

Proof: Choose an ordered basis $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ for E_λ and extend it to an ordered basis $\{\vec{v}_1, \dots, \vec{v}_p, \vec{v}_{p+1}, \dots, \vec{v}_n\}$ for V . \curvearrowleft p = \dim(E_\lambda)

Then: $[T]_\beta = \begin{pmatrix} | & | \\ [T(\vec{v}_1)]_\beta & [T(\vec{v}_p)]_\beta \\ | & | \end{pmatrix} = \begin{pmatrix} \lambda & 0 & 0 & \cdots & 0 \\ 0 & \lambda & 0 & \cdots & 0 \\ 0 & 0 & \lambda & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix} \boxed{\text{diag}}$

$$= \boxed{\begin{array}{c|c} \lambda I_p & B \\ \hline 0 & C \end{array}}$$

$$\begin{aligned}
 \Rightarrow f_T(t) &= \det \left(\begin{array}{c|c} (\lambda-t)I_p & B \\ \hline 0 & C-tI_{n-p} \end{array} \right) \\
 &= \det((\lambda-t)I_p) \underline{\det(C-tI_{n-p})} \\
 &= (\lambda-t)^p \det(C-tI_{n-p}) \\
 \therefore (\lambda-t)^p &\mid f_T(t) \\
 \therefore \mu_T(\lambda) &\geq p = \gamma_T(\lambda)
 \end{aligned}$$

Lemma: Let T be a linear operator, and let $\lambda_1, \lambda_2, \dots, \lambda_k$ distinct eigenvalues of T . For each $i=1, 2, \dots, k$, let $\vec{v}_i \in E_{\lambda_i}$.


If $\vec{v}_1 + \vec{v}_2 + \dots + \vec{v}_k = \vec{0}$, then $\vec{v}_i = \vec{0}$ for all i .

Proof: If not, say $\vec{v}_1, \dots, \vec{v}_s \neq \vec{0}$

then:

$$\vec{v}_1 + \vec{v}_2 + \dots + \vec{v}_s = \vec{0}$$

It contradicts to our previous proposition that $\vec{v}_1, \dots, \vec{v}_s$ must be lin. independent.



Proposition: Let T be a linear operator, and let $\lambda_1, \lambda_2, \dots, \lambda_k$ be distinct eigenvalues of T . For each $i=1, 2, \dots, k$, let $S_i \subset E_{\lambda_i}$ be a finite linearly independent subset. Then:

$S = S_1 \cup S_2 \cup \dots \cup S_k$ is a linearly independent subset of V .

Proof: Write $S_i = \{\vec{v}_{i1}, \vec{v}_{i2}, \dots, \vec{v}_{in_i}\}$ for $i=1, 2, \dots, k$. Suppose $\exists a_{ij} \in F$ for $1 \leq j \leq n_i$ and $1 \leq i \leq k$ such that

$$\sum_{i=1}^k \sum_{j=1}^{n_i} a_{ij} \vec{v}_{ij} = \vec{0}$$

Then: $\begin{matrix} \vec{v}_1 & \vec{v}_2 & \dots & \vec{v}_k \\ \downarrow & \downarrow & \dots & \downarrow \\ w_1 + w_2 + \dots + w_k & = \vec{0} & \in E_{\lambda_1} \end{matrix}$

$$w_i = \sum_{j=1}^{n_i} a_{ij} \vec{v}_{ij} = \vec{0} \quad \text{for all } i.$$

Then: $a_{ij} = 0$ for all i and j
(for S_i are lin. independent for all i .
i. $S_1 \cup S_2 \cup \dots \cup S_k$ is linearly independent.

Theorem: Let T be a linear operator on a finite dimensional vector space V such that the characteristic polynomial splits. Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be distinct eigenvalues of T .

Then: (a) T is diagonalizable iff: $M_T(\lambda_i) = g_T(\lambda_i)$
for $i=1, 2, \dots, k$

(b) If T is diagonalizable and β_i is an ordered basis for E_{λ_i} for each i , then $\beta := \beta_1 \cup \beta_2 \cup \dots \cup \beta_k$ is an ordered basis for V consisting of eigenvectors.

(so that $[T]_{\beta}$ is a diagonal matrix)

Proof: Write $n = \dim(V)$, and $m_i = M_T(\lambda_i)$ and $d_i = \dim(E_{\lambda_i})$ for all i . $\dim(E_{\lambda_i})$

Suppose T is diagonalizable and β is a basis for V consisting of eigenvectors of T .

(e.g. $\beta = \{\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4, \vec{v}_5, \dots, \vec{v}_n\}$)

The diagram illustrates the decomposition of the basis β into eigenspaces. The vectors \vec{v}_1 and \vec{v}_3 are highlighted in yellow, while \vec{v}_2 , \vec{v}_4 , and \vec{v}_5 are highlighted in pink. Green arrows connect the first three vectors to a blue oval labeled E_{λ_1} , indicating they are eigenvectors corresponding to the eigenvalue λ_1 . Similarly, the last three vectors connect to a green oval labeled E_{λ_j} , indicating they are eigenvectors corresponding to the eigenvalue λ_j .

For each i , let $\beta_i = \beta \cap E_{\lambda_i}$ and $n_i \stackrel{\text{def}}{=} \#\beta_i$

Then: $n_i \leq d_i = \dim(E_{\lambda_i})$ ($\because \beta_i$ is lin. independent)

Also, $d_i \leq m_i$ (last lecture)

So, we have $n_i \leq d_i \leq m_i$ for all i .

$$\therefore n = \sum_{i=1}^k n_i \leq \sum_{i=1}^k d_i \leq \sum_{i=1}^k m_i = n = \dim(V)$$

$$\therefore \sum_{i=1}^k d_i - \sum_{i=1}^k n_i = 0 \Leftrightarrow \sum_{i=1}^k (d_i - n_i) = 0$$
$$\Rightarrow d_i = n_i \text{ for all } i.$$

$$\therefore \sum_{i=1}^k m_i - \sum_{i=1}^k d_i = 0 \Leftrightarrow \sum_{i=1}^k (m_i - d_i) = 0$$
$$\Rightarrow d_i = m_i \text{ for all } i.$$

$$\therefore n_i = d_i = m_i \text{ for all } i$$

(So, β_i is a basis of E_{n_i})

Conversely, suppose $m_i = d_i \forall i$.

For each i , let β_i be the ordered basis of E_{λ_i} and let $\beta = \beta_1 \cup \beta_2 \cup \dots \cup \beta_k$.


Then: from previous proposition, we know β is linearly independent.

$$\text{But } \# \beta = \sum_{i=1}^k d_i = \sum_{i=1}^k m_i = n = \dim(V)$$

$$\begin{matrix} |\beta_1| + |\beta_2| + \dots + |\beta_k| \\ \text{dim}(E_{\lambda_1}) \quad \text{dim}(E_{\lambda_2}) \quad \text{dim}(E_{\lambda_k}) \\ d_1 \end{matrix}$$

$\therefore \beta$ is a basis for V of eigenvectors

$\therefore T$ is diagonalizable.



If S_1, \dots, S_k are L.I.
then $S_1 \cup S_2 \cup \dots \cup S_k$ is L.I. ??