

Thm If $f, g: A \xrightarrow{C\mathbb{R}^n} \mathbb{R}$ are continuous at $\vec{a} \in A$, then

(1) $f(\vec{x}) \pm g(\vec{x})$, $kf(\vec{x})$, $f(\vec{x})g(\vec{x})$ are continuous at \vec{a} ,
where k is a constant.

(2) $\frac{f(\vec{x})}{g(\vec{x})}$ is continuous at \vec{a} provided $g(\vec{a}) \neq 0$

Consequences:

(i) All polynomials of multi-variables are continuous (on \mathbb{R}^n)


(ii) All rational functions of multi-variables are continuous
on their domain of definition

(rational function def $\frac{P(\vec{x})}{Q(\vec{x})}$ for some polynomials $P(\vec{x})$ & $Q(\vec{x})$)
"domain of definition" of $\frac{P(\vec{x})}{Q(\vec{x})} = \mathbb{R}^n \setminus \{ \vec{x} : Q(\vec{x}) = 0 \}$

egs (1) $x^3 + 3yz + z^2 - x + 7y$ is a polynomial on \mathbb{R}^3
& is continuous on \mathbb{R}^3

(2) $\frac{x^3 + y^2 + 4z}{x^2 + y^2}$ is a rational function on \mathbb{R}^3

domain of definition = $\mathbb{R}^3 \setminus \{ (0, 0, z) \}$
= $\mathbb{R}^3 \setminus \{ z\text{-axis} \}$



& is continuous on $\mathbb{R}^3 \setminus \{ z\text{-axis} \}$.

Fact: Let \vec{a} be a zero of polynomial $Q(\vec{x})$ (ie. $Q(\vec{a})=0$)

Then the rational function $r(\vec{x}) = \frac{P(\vec{x})}{Q(\vec{x})}$ can be

"extended to a function continuous at $\vec{a} \iff \lim_{\vec{x} \rightarrow \vec{a}} r(\vec{x})$ exists".

egs (1) $f(x,y) = \frac{xy+y^3}{x^2+y^2}$ (in \mathbb{R}^2)

Note $x^2+y^2=0 \iff (x,y)=(0,0)$

\therefore Domain of definition of f is $\mathbb{R}^2 \setminus \{(0,0)\}$

$$\lim_{\substack{(x,y) \rightarrow (0,0) \\ y=mx}} f(x,y) = \lim_{\substack{(x,y) \rightarrow (0,0) \\ y=mx}} \frac{xy+y^3}{x^2+y^2} = \lim_{x \rightarrow 0} \frac{x(mx) + (mx)^3}{x^2 + (mx)^2}$$

$$= \lim_{x \rightarrow 0} \frac{m+mx}{1+m^2} = \frac{m}{1+m^2}$$

Different limits in different directions

$$\Rightarrow \lim_{(x,y) \rightarrow (0,0)} f(x,y) \text{ DNE}$$

$\therefore \frac{xy+y^3}{x^2+y^2}$ cannot be extended to a function cts. at $(0,0)$. ~~##~~

(2) $g(x,y) = \frac{x^4-y^4}{x^2+y^2}$ (Note $x^2+y^2=0 \iff (x,y)=(0,0)$)

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^4-y^4}{x^2+y^2} = \lim_{r \rightarrow 0} \frac{r^4(\cos^4\theta - \sin^4\theta)}{r^2} = \lim_{r \rightarrow 0} r^2(\cos^4\theta - \sin^4\theta) = 0$$

(Squeeze Thm)

$\therefore g(x,y) = \frac{x^4 - y^4}{x^2 + y^2}$ can be extended to a function cts. at $(0,0)$,

In fact $g(x,y) = \begin{cases} \frac{x^4 - y^4}{x^2 + y^2} & \text{if } (x,y) \neq (0,0) \\ 0 & \text{if } (x,y) = (0,0) \end{cases} \quad \left(= x^2 - y^2 \right)$
is the required extension.

Thm If $\bullet f: A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous at $\vec{a} \in A$,

$\bullet g(x)$ is a 1-variable function continuous at $f(\vec{a})$

Then $g \circ f(\vec{x})$ (def $g(f(\vec{x}))$) is continuous at \vec{a} , and

$$\lim_{\vec{x} \rightarrow \vec{a}} g(f(\vec{x})) = g\left(\lim_{\vec{x} \rightarrow \vec{a}} f(\vec{x})\right) = g(f(\vec{a})).$$

egs: (1) x_k k -th coordinate function are continuous, $\forall k=1, \dots, n$

$\left\{ \begin{array}{l} \bullet |x| \text{ is also continuous} \Rightarrow |x_k| \text{ are continuous.} \\ \bullet \ln|x| \text{ is continuous for } |x| > 0 \Rightarrow \ln|x_k| \text{ are continuous} \\ \text{if } |x_k| > 0. \end{array} \right.$

(2) $\sin(x^2 + y^2)$, e^{x-y} , $\cos\left(\frac{1}{x^2 + y^2}\right)$ (except $(x,y) = (0,0)$)

$r = \sqrt{x^2 + y^2}$ are continuous on their domains.

Partial Derivatives

Def: Let $\bullet \Omega \subseteq \mathbb{R}^n$ be open

$\bullet f: \Omega \rightarrow \mathbb{R}$ be a function

Then the i -th partial derivative of f at $\vec{x} = (x_1, \dots, x_n) \in \Omega$ is defined by

$$\frac{\partial f}{\partial x_i}(\vec{x}) = \frac{\partial f}{\partial x_i}(x_1, \dots, x_n)$$

$$= \lim_{h \rightarrow 0} \frac{f(x_1, \dots, x_i+h, \dots, x_n) - f(x_1, \dots, x_i, \dots, x_n)}{h}$$

(provided the limit exists)

Remarks: (1) Ω open "ensures" $(x_1, \dots, x_i+h, \dots, x_n) \in \Omega$ for small h so that $f(x_1, \dots, x_i+h, \dots, x_n)$ is defined.

(2) If $n=1$, $\frac{\partial f}{\partial x} = \frac{df}{dx}$

(3) If $n=2$, we usually write

$$\frac{\partial f}{\partial x}(x, y) = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h}$$

$$\frac{\partial f}{\partial y}(x, y) = \lim_{h \rightarrow 0} \frac{f(x, y+h) - f(x, y)}{h}$$

(4) In practice, $\frac{\partial f}{\partial x_i}$ can be calculated as derivative in one variable x_i by regarding other variables as constants.

(5) Other notations: $\frac{\partial f}{\partial x} = \partial_1 f = D_1 f = \nabla_1 f = f_x = f_1$
 $\frac{\partial f}{\partial y} = \partial_2 f = D_2 f = \nabla_2 f = f_y = f_2$

eg $f(x,y) = x^2 + y^2$

$$\frac{\partial f}{\partial x} = \frac{\partial}{\partial x} (x^2 + y^2) = 2x + 0 = 2x$$

↑ regarded as constant

$$\frac{\partial f}{\partial y} = \frac{\partial}{\partial y} (x^2 + y^2) = 0 + 2y = 2y$$

$\frac{\partial y^2}{\partial x} = 0$
 $\frac{\partial x^2}{\partial y} = 0$

Note: As the point (1, -1)


$$\frac{\partial f}{\partial x}(1, -1) = 2 \quad \& \quad \frac{\partial f}{\partial y}(1, -1) = -2$$

$\downarrow 0$ $\uparrow 0$

f increases as x increases at (1, -1)

f decreases as y increases at (1, -1)

$$f(x,y) = x^2 + y^2 = (\text{dist. to } (0,0))^2$$



eg: $f(x, y, z) = xy^2 - \cos(xz)$


$$f_x = \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} (xy^2 - \cos(xz)) = y^2 + z \sin(xz)$$

$$f_y = \frac{\partial f}{\partial y} = \frac{\partial}{\partial y} (xy^2 - \cos(xz)) = 2xy$$

$$f_z = \frac{\partial f}{\partial z} = \frac{\partial}{\partial z} (xy^2 - \cos(xz)) = x \sin(xz) \quad \text{✗}$$

eg $f(x, y) = \begin{cases} 1, & \text{if } xy \geq 0 \\ 0, & \text{if } xy < 0 \end{cases}$

Find $\frac{\partial f}{\partial x}(1, 1)$, $\frac{\partial f}{\partial x}(0, 0)$, & $\frac{\partial f}{\partial x}(0, 1)$



Solu: For (1, 1)

$$\lim_{h \rightarrow 0} \frac{f(1+h, 1) - f(1, 1)}{h}$$

(small |h|)

$$= \lim_{h \rightarrow 0} \frac{1-1}{h} = 0$$

$$\therefore \frac{\partial f}{\partial x}(1, 1) = 0$$

For (0, 0)

$$\frac{\partial f}{\partial x}(0, 0) = \lim_{h \rightarrow 0} \frac{f(0+h, 0) - f(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{1-1}{h} = 0$$

For (0, 1),

$$\lim_{\substack{h \rightarrow 0 \\ h > 0}} \frac{f(0+h, 1)}{h} = \lim_{h \rightarrow 0} \frac{1-1}{h} = 0$$

$$\lim_{\substack{h \rightarrow 0 \\ h < 0}} \frac{f(0+h, 1)}{h} = \lim_{h \rightarrow 0} \frac{0-1}{h} \quad \text{DNE}$$

$$\therefore \frac{\partial f}{\partial x}(0, 1) \text{ DNE} \quad \#$$

(Note: $\frac{\partial f}{\partial x}(0, 0)$ exists, but f is not continuous at $(0, 0)$)
check: $\frac{\partial f}{\partial y}(0, 0) = 0$ (Optional Ex.)