

## Tutorial 5

Exercises 13.2

Q39 At what points  $(x, y, z)$  in space are the functions continuous?

a.  $h(x, y, z) = \ln(z - x^2 - y^2 - 1)$

b.  $h(x, y, z) = \frac{1}{z - \sqrt{x^2 + y^2}}$

a.  $h$  is continuous at  $(x, y, z)$  if  $z - x^2 - y^2 - 1 > 0$ .  
 $z > x^2 + y^2 + 1$ .

$\therefore h$  is continuous at all  $(x, y, z)$  such that  $z > x^2 + y^2 + 1$ .

b.  $h$  is continuous at  $(x, y, z)$  if  $z - \sqrt{x^2 + y^2} \neq 0$ .  
 $z \neq \sqrt{x^2 + y^2}$

$\therefore h$  is continuous at all  $(x, y, z)$  such that  $z \neq \sqrt{x^2 + y^2}$ .

Q59 Does knowing that  $1 - \frac{x^2 y^2}{3} < \frac{\tan^{-1} xy}{xy} < 1$

tell you anything about  $\lim_{(x,y) \rightarrow (0,0)} \frac{\tan^{-1} xy}{xy}$ ?

Give reasons for your answer.

Squeeze Theorem (Sandwich Theorem)

Let  $f, g, h : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$  be functions of  $n$ -variables.

If  $\begin{cases} g(\vec{x}) \leq f(\vec{x}) \leq h(\vec{x}) \text{ near } \vec{a} \in \Omega \text{ and} \\ \lim_{\vec{x} \rightarrow \vec{a}} g(\vec{x}) = \lim_{\vec{x} \rightarrow \vec{a}} h(\vec{x}) = L. \end{cases}$

Then  $\lim_{\vec{x} \rightarrow \vec{a}} f(\vec{x}) = L$

$$\lim_{(x,y) \rightarrow (0,0)} \left(1 - \frac{x^2 y^2}{3}\right) = 1, \quad \lim_{(x,y) \rightarrow (0,0)} (1) = 1$$

$\Rightarrow \lim_{(x,y) \rightarrow (0,0)} \frac{\tan xy}{xy} = 1$  by the Squeeze Theorem.

Q71 Define  $f(0,0)$  in a way that extends  $f$  to be continuous at the origin.

$$f(x,y) = \ln\left(\frac{3x^2 - x^2 y^2 + 3y^2}{x^2 + y^2}\right)$$

f is continuous at  $(0,0)$  if  $\lim_{(x,y) \rightarrow (0,0)} f(x,y) = f(0,0)$ .

$$\lim_{(x,y) \rightarrow (0,0)} f(x,y) = \lim_{(x,y) \rightarrow (0,0)} \ln\left(\frac{3x^2 - x^2 y^2 + 3y^2}{x^2 + y^2}\right)$$

$$= \lim_{(x,y) \rightarrow (0,0)} \ln\left(\frac{3(x^2 + y^2) - x^2 y^2}{x^2 + y^2}\right)$$

$$= \lim_{r \rightarrow 0} \ln\left(\frac{3r^2 - r^4 \cos^2 \theta \sin^2 \theta}{r^2}\right)$$

$$= \lim_{r \rightarrow 0} \ln(3 - r^2 \cos^2 \theta \sin^2 \theta)$$

$$= \ln 3$$

$$\text{Define } f(0,0) = \ln 3$$

### Exercises 13.3

#### Partial Derivatives

$$\frac{\partial f}{\partial x}(x,y) = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h}$$

(regarding  $y$  as constant)

$$\frac{\partial f}{\partial y}(x,y) = \lim_{h \rightarrow 0} \frac{f(x, y+h) - f(x, y)}{h}$$

(regarding  $x$  as constant)

Polar coordinates :  
 $x = r \cos \theta$   
 $y = r \sin \theta$   
 $r^2 = x^2 + y^2$

Q21  $f(x, y) = \int_x^y g(t) dt$  (g continuous for all t)

Find  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$ .

$$\begin{aligned}\frac{\partial f}{\partial x} &= \frac{\partial}{\partial x} \left( \int_x^y g(t) dt \right) = \frac{\partial}{\partial x} \left( - \int_y^x g(t) dt \right) \\ &= -g(x)\end{aligned}$$

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y} \left( \int_x^y g(t) dt \right) \\ &\quad \text{regarding } x \text{ as constant} \\ &= g(y)\end{aligned}$$

Fundamental  
Theorem of Calculus

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

↑  
constant

Q47  $w = x^2 \tanh(xy)$

Find all the second-order partial derivatives of the function.

$$\left(\text{treat } y \text{ constant}\right) \frac{\partial w}{\partial x} = x^2 \sec^2(xy) \cdot y + 2x \tan(xy) = x^2 y \sec^2(xy) + 2x \tan(xy)$$

$$\left(\text{treat } x \text{ constant}\right) \frac{\partial w}{\partial y} = x^2 \sec^2(xy) \cdot x = x^3 \sec^2(xy)$$

$$\begin{aligned}\frac{\partial^2 w}{\partial x^2} &= y (x^2 (2 \sec(xy) \sec(xy) \tan(xy) \cdot y) + 2x \sec^2(xy)) \\ &\quad + 2x \sec^2(xy) \cdot y + 2 \tan(xy) \\ &= 2x^2 y^2 \sec^2(xy) \tan(xy) + 4xy \sec^2(xy) + 2 \tan(xy)\end{aligned}$$

$$\begin{aligned}\frac{\partial^2 w}{\partial y^2} &= x^3 2 \sec(xy) \cdot \sec(xy) \tan(xy) \cdot x \\ &= 2x^4 \sec^2(xy) \tan(xy)\end{aligned}$$

$$\frac{\partial^2 w}{\partial y \partial x} = 2x^3 y \sec^2(xy) \tan(xy) + 3x^2 \sec^2(xy)$$

$$\frac{\partial^2 w}{\partial x \partial y} = \frac{\partial}{\partial x} (x^3 \sec^2(xy)) = x^3 \cdot 2 \sec(xy) \sec(xy) \tan(xy) \cdot y + 3x^2 \sec^2(xy)$$