

# MATH 3310 Assignment 3

Due: November 5, 2024

1. Consider the following linear system  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 3 \end{pmatrix} \text{ and } b = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$$

- (a) Determine whether the Jacobi method converges.
- (b) Using initial approximation  $x^0 = (3, 2, 1)^T$ , conduct the first two Jacobi iterations.
- (c) Determine whether the Gauss-Seidel method converges.
- (d) Using initial approximation  $x^0 = (2, 1, 0)^T$ , conduct the first two Gauss-Seidel iterations.

2. Consider the following linear system  $Ax = b$ , where

$$A = \begin{pmatrix} 2 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 1 & 3 \end{pmatrix} \text{ and } b = \begin{pmatrix} 8 \\ 8 \\ 8 \end{pmatrix}$$

- (a) Determine whether the SOR method converges if  $\omega = 1.5$ .
- (b) Using initial approximation  $x^{(0)} = (6, 4, 0)^T$ , conduct the first two SOR iteration.

3. Consider solving  $Ax = b$  with the following iterative scheme.

$$x_{k+1} = x_k + \alpha(b - Ax_k)$$

where  $\alpha > 0$ . Assume all eigenvalues  $\lambda_i$  of  $A$  are real, and such that  $\lambda_{min} \leq \lambda_i \leq \lambda_{max}$ ,  $\lambda_{min}, \lambda_{max}$  are the smallest and largest eigenvalues of  $A$ , respectively.

- (a) If  $\lambda_{min} < 0$  and  $\lambda_{max} > 0$ , prove that this scheme always diverges for some initial guess.
- (b) Assume  $\lambda_{min} > 0$ , what are the sufficient and necessary conditions on  $\alpha$  for the iterative scheme to converge?

- (c) Assume  $\lambda_{min} > 0$ , what is the best value  $\alpha_{opt}$  for  $\alpha$ , i.e., the value of  $\alpha$  minimizing the convergence factor? And, what is the convergence factor in such case?

4. Consider an  $n \times n$  tridiagonal matrix of the form

$$T_\alpha = \begin{pmatrix} \alpha & -1 & & & \\ -1 & \alpha & -1 & & \\ & -1 & \ddots & \ddots & \\ & & \ddots & \ddots & -1 \\ & & & -1 & \alpha \end{pmatrix},$$

where  $\alpha$  is a real parameter.

- (a) Verify that the eigenvalues of  $T_\alpha$  are given by

$$\lambda_j = \alpha - 2 \cos(j\theta), \quad j = 1, \dots, n,$$

where

$$\theta = \frac{\pi}{n+1},$$

and that an eigenvector associated with each  $\lambda_j$  is

$$q_j = [\sin(j\theta), \sin(2j\theta), \dots, \sin(nj\theta)]^T.$$

Under what condition on  $\alpha$  does this matrix become positive definite?

- (b) Let  $\alpha = 2$ .
- Will the Jacobi iteration converge for this matrix? If so, what will its convergence factor be?
  - Will the Gauss-Seidel iteration converge for this matrix?
  - For which values of  $\omega$  will the SOR iteration converge?

5. Let  $B$  be a matrix with the following structure:

$$B = \begin{pmatrix} O & B_{12} \\ B_{21} & O \end{pmatrix},$$

and let  $L$  and  $U$  be the lower and upper triangular parts of  $B$ , respectively.

- (a) If  $\mu$  is an eigenvalue of  $B$ , then so is  $-\mu$ .
- (b) The eigenvalues of the matrix

$$B(\alpha) = \alpha L + \frac{1}{\alpha} U$$

defined for  $\alpha \neq 0$  are independent of  $\alpha$ .