



Topics in Numerical Analysis II

Computational Inverse Problems

Lecturer: Bangti Jin (b.jin@cuhk.edu.hk)

Chinese University of Hong Kong

September 23, 2024



Outline

1 Tikhonov regularization - Nonlinear case



Review of previous lecture

Tikhonov regularization for linear inverse problems

$$x_\alpha = \arg \min_x \|Ax - y\|^2 + \alpha \|x\|^2$$

key idea: restrict the candidate solutions to a “compact” subset

- the method is well defined: x_α exists and is unique, expressed by

$$x_\alpha = \sum_i \frac{s_i}{s_i^2 + \alpha} (y, u_i) v_i$$

- it is stable under perturbation of α and y
- it is consistent under a prior choice

$$\lim_{\delta \rightarrow 0^+} \alpha(\delta)^{-1} \delta^2 = 0 \quad \text{and} \quad \lim_{\delta \rightarrow 0^+} \alpha(\delta) = 0$$

- for $x^\dagger \in \text{range}(A^*)$, it has a convergence rate $O(\delta^{\frac{1}{2}})$



Morozov's discrepancy principle: given noisy data y^δ with $\|y^\delta - y^\dagger\| = \delta$, choose α s.t.

$$\|Ax_\alpha^\delta - y^\delta\| = \delta$$

- $\alpha(\delta)$ is uniquely defined under minor conditions
- Tikhonov + Morozov is convergent
- Tikhonov + Morozov + source condition has convergence rate



(generalized) Tikhonov regularization

$$J_{\alpha}(x) = \|Ax - y\|^2 + \alpha \|L(x - x_0)\|^2$$

- L : differential operator
- intuition: look for a solution with small derivative $\|L(x - x_0)\| \Rightarrow$ weak compactness ...
- the penalty term often induces a (weak) compact subset in Sobolev spaces





The implementation of Tikhonov regularization is as easy as for the standard case. If $X = \mathbb{R}^n$ and $Y = \mathbb{R}^m$, the operator $L \in \mathbb{R}^{\ell \times n}$ and the Tikhonov functional can be given as



$$J_{\alpha}(x) = \|Kx - z\|^2$$


with

$$K = \begin{bmatrix} A \\ \sqrt{\alpha}L \end{bmatrix} \quad \text{and} \quad z = \begin{bmatrix} y \\ \sqrt{\alpha}Lx_0 \end{bmatrix}$$


If L is properly chosen, then all singular values of K are positive and the regularized solution x_{α} can be computed by solving the least squares problem.







H^1 seminorm



H^2 seminorm





Nonlinear inverse problems

$$F(x) = y,$$

with $F : X \rightarrow Y$ being nonlinear operator (possibly compact) between Hilbert spaces X and Y

- (i) F is continuous
- (ii) F is **weakly sequentially closed**, i.e., $(x_n) \subset D(F)$, weak convergence of x_n to $x^* \in X$ and weak convergence of $F(x_n)$ to y^* in Y imply $y^* = F(x^*) \in Y$

model nonlinear inverse problem: electrical impedance tomography




© Heinrich et al, Intensive Care Med., 2006.

mathematical model

$$\begin{cases} -\nabla \cdot (\sigma \nabla u) = 0, & \text{in } \Omega \\ \sigma \partial_\nu u = f, & \text{on } \partial\Omega \end{cases}$$

with **many pairs** of boundary data (Neumann to Dirichlet map)
goal: to recover the conductivity σ from DtN map

model nonlinear inverse problem: optical tomography blood volume / oxygen saturation of infant



© J. C. Hebden, T. Austin, Eur Radiol 2007


mathematical model (radiative transfer equation)

$$\mathbf{v} \cdot \nabla_{\mathbf{x}} u + \sigma(\mathbf{x})u = \int_{\mathbb{S}} k(\mathbf{x}, \mathbf{v}, \mathbf{v}') u(\mathbf{x}, \mathbf{v}') d\mathbf{v}',$$

goal: recover scattering kernel k and absorption coefficient σ



inverse scattering problems



with incident wave $u^i = e^{ikx \cdot d}$, $u = u^i + u^s$ satisfies,

$$\begin{cases} -\Delta u - k^2 n(x)^2 u = 0, & \text{in } \mathbb{R}^d, \\ \lim_{r \rightarrow \infty} r^{\frac{d-1}{2}} \left(\frac{\partial u^s}{\partial r} - iku^s \right) = 0 & r = |x|. \end{cases}$$

applications: remote sensing, radar



Tikhonov regularization

$$J_{\alpha}(x) = \|F(x) - y\|^2 + \alpha\|x\|^2$$

and approximation x_{α}

$$x_{\alpha} \in \arg \min_{x \in D(F)} J_{\alpha}(x)$$



x_0 -minimum norm solution $x^\dagger$

$$\|F(x^\dagger) - y^\dagger\| = \min\{\|F(x) - y^\dagger\| : x \in D(F)\}$$

and the set is denoted by S , and

$$\|x^\dagger - x_0\| = \min_{x \in S} \|x - x_0\|$$

- generally the existence and uniqueness of $x^\dagger$ is nontrivial
- existence of x_0 -minimum norm solution
- the choice of x_0 is crucial in practice



existence of a minimizer (by compactness argument)

- $J_\alpha(x) \geq 0 \Rightarrow \exists$ a minimizing sequence $(x^n) \subset D(F)$

$$\lim_{n \rightarrow \infty} J_\alpha(x^n) = \inf_{x \in D(F)} J_\alpha(x)$$

- The sequence (x^n) is uniformly bdd. in X
The sequence $(F(x^n))$ is uniformly bdd. in Y
- $\exists$ a subsequence (x^{n_k}) converging to $x^* \in D(F)$ weakly in X ,
and $(F(x^{n_k}))$ converges weakly to $F(x^*)$ (by weak seq.
closedness of F)
- weak lower semi-continuity of norm $\Rightarrow \exists$ a minimizer

the whole argument holds only on a subsequence



stability of the x_α

Let $\alpha > 0$, and (y^k) and (x^k) be seq. with $y^k \rightarrow y^\delta$ and x^k be a minimizer of J_α (for y^k). Then there exists a convergent subseq. of (x^k) and the limit of every convergent subseq. is a minimizer of J_α .



The preceding argument implies that x^* is a minimizer of J_α , and

$$\lim_{k \rightarrow \infty} \|F(x^k) - y^k\|^2 + \alpha \|x^k - x_0\|^2 = \|F(x^*) - y^\delta\|^2 + \alpha \|x^* - x_0\|^2.$$

If $x^k \not\rightarrow x^*$, then

$$c := \limsup \|x^k - x_0\| > \|x^* - x_0\|$$

and there exists a subseq. of (x^k) s.t. $x^{k_j} \rightharpoonup x^*$, $F(x^{k_j}) \rightharpoonup F(x^*)$ and $\lim_{k \rightarrow \infty} \|x^{k_j} - x_0\| = c$. Thus we obtain

$$\begin{aligned} \lim_{j \rightarrow \infty} \|F(x^{k_j}) - y^{k_j}\|^2 &= \|F(x^*) - y^\delta\|^2 + \alpha(\|x^* - x_0\|^2 - \lim_{j \rightarrow \infty} \|x^{k_j} - x_0\|^2) \\ &= \|F(x^*) - y^\delta\|^2 + \alpha(\|x^* - x_0\|^2 - c^2) < \|F(x^*) - y^\delta\|^2 \end{aligned}$$

contradicts the weak lower semicontinuity. This shows $x^k \rightarrow x^*$.



convergence:

$(y^\delta)_{\delta>0}$ be seq. with $\delta = \|y^\delta - y^\dagger\|$ and $x_{\alpha(\delta)}^\delta$ be a minimizer of $J_{\alpha(\delta)}$ (for y^δ). If

$$\lim_{\delta \rightarrow 0^+} \alpha(\delta) = \lim_{\delta \rightarrow 0^+} \alpha(\delta)^{-1} \delta^2 = 0$$

then there exists a convergent subseq. of $(x_{\alpha(\delta)}^\delta)$ and the limit of every convergent subseq. is an x_0 -minimum norm solution.



convergence rate analysis

Let $D(F)$ be **convex**, $y^\delta \in Y$ with $\|y^\delta - y^\dagger\| \leq \delta$, and $x^\dagger$ be an x_0 -minimum norm solution. Moreover, the following condition holds

- (i) F is **Frechet differentiable**.
- (ii) $\exists L > 0$ s.t. $\|F'(x^\dagger) - F'(z)\| \leq L\|x^\dagger - z\|$ for all $z \in D(F)$
- (iii) $\exists w \in Y$ s.t. $x^\dagger - x_0 = F'(x^\dagger)^* w$ and $L\|w\| < 1$.

Then with $\alpha \sim O(\delta)$, we have

$$\|x_\alpha^\delta - x^\dagger\| = O(\delta^{\frac{1}{2}}).$$



■ minimizing property (with $x_0 = 0$)

$$\|F(x_\alpha^\delta) - y^\delta\|^2 + \alpha\|x_\alpha^\delta\|^2 \leq \|F(x^\dagger) - y^\delta\|^2 + \alpha\|x^\dagger\|^2$$

■ completing squares

$$\begin{aligned} & \|F(x_\alpha^\delta) - y^\delta\|^2 + \alpha\|x_\alpha^\delta - x^\dagger\|^2 \\ & \leq \|F(x^\dagger) - y^\delta\|^2 - 2\alpha\langle x^\dagger, x_\alpha^\delta - x^\dagger \rangle \\ & = \|F(x^\dagger) - y^\delta\|^2 - 2\alpha\langle F'(x^\dagger)^* w, x_\alpha^\delta - x^\dagger \rangle \\ & = \|F(x^\dagger) - y^\delta\|^2 - 2\alpha\langle w, F'(x^\dagger)(x_\alpha^\delta - x^\dagger) \rangle \end{aligned}$$



■ Lipschitz continuity of the derivative:

$$\|F(x_\alpha^\delta) - F(x^\dagger) - F'(x^\dagger)(x_\alpha^\delta - x^\dagger)\| \leq \frac{L}{2} \|x_\alpha^\delta - x^\dagger\|^2,$$

i.e.,

$$\|F'(x^\dagger)(x_\alpha^\delta - x^\dagger)\| \leq \|F(x_\alpha^\delta) - F(x^\dagger)\| + \frac{L}{2} \|x_\alpha^\delta - x^\dagger\|^2$$

■ triangle inequality + Cauchy-Schwarz inequality

$$\begin{aligned} & \|F(x_\alpha^\delta) - y^\delta\|^2 + \alpha(1 - L\|w\|)\|x_\alpha^\delta - x^\dagger\|^2 \\ & \leq \|F(x^\dagger) - y^\delta\|^2 + \alpha\|w\|\|F(x_\alpha^\delta) - y^\delta\| + \alpha\|w\|\|y^\delta - F(x^\dagger)\| \end{aligned}$$

■ Young inequality

$$\|x_\alpha^\delta - x^\dagger\| \leq c\sqrt{\delta}$$

discrepancy principle:

$$\|F(x_\alpha^\delta) - y^\delta\| = \delta$$

- under certain conditions, there exists $\alpha(\delta)$ for DP, if δ is sufficiently small
- $x_{\alpha(\delta)}^\delta \rightarrow x^\dagger$ as $\delta \rightarrow 0$, if $x^\dagger$ is unique
- under the conditions of the theorem,

$$\|x_{\alpha(\delta)}^\delta - x^\dagger\|^2 \leq 4(1 - L\|w\|)^{-1}\|w\|\delta$$



Model problem

Most inverse problems for PDEs are nonlinear in nature
model problem:

$$-\Delta u + qu = f \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

with given $q \in L^\infty(\Omega)$, $q \geq 0$, and $f \in L^2(\Omega)$

The forward is well-posed: there exists a unique solution $u \in H_0^1(\Omega)$.

Lax-Milgram theorem If $a(\cdot, \cdot) : V \times V \rightarrow \mathbb{R}$ is a **continuous and coercive** bilinear form on V and ℓ is a bounded linear functional on V , then the problem

$$a(u, v) = \ell(v) \quad \forall v \in V$$

has a unique solution $u \in V$.



multiply the equation with $v \in H_0^1(\Omega)$ and integration by parts

$$\underbrace{\int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\Omega} q u v dx}_{a(u,v)} = \underbrace{\int_{\Omega} f v dx}_{\ell(v)}$$

continuity & coercivity of $a(\cdot, \cdot)$

$$\begin{aligned} |a(u, v)| &\leq \int_{\Omega} |\nabla u| |\nabla v| dx + \int_{\Omega} q |u| |v| dx \\ &\leq \|\nabla u\|_{L^2(\Omega)} \|\nabla v\|_{L^2(\Omega)} + \|q\|_{L^\infty(\Omega)} \|u\|_{L^2(\Omega)} \|v\|_{L^2(\Omega)} \\ &\leq c \|u\|_{H^1(\Omega)} \|v\|_{H^1(\Omega)} \\ a(u, u) &\geq \|\nabla u\|_{L^2(\Omega)}^2 \geq c \|u\|_{H^1(\Omega)}^2 \quad (\text{Poincare inequality}) \end{aligned}$$

boundedness of ℓ (linearity...)

$$|\ell(v)| \leq \int_{\Omega} |f| |v| dx \leq \|f\|_{L^2(\Omega)} \|v\|_{L^2(\Omega)} \leq \|f\|_{L^2(\Omega)} \|v\|_{H^1(\Omega)}$$

Lax-Milgram theorem $\Rightarrow \exists! u \in H_0^1(\Omega)$ (for any given q)



Inverse problem: given $g \approx u^\delta$ in Ω , find q .

How bad is the inverse problem:

$$q = \frac{f + \Delta u}{u} \quad \text{twice differentiation in space}$$

(numerical differential is important and relatively easy !)
so the inverse problem is not that hard (in theory) ...



ill-posedness by comparing the solution for two differential potentials:

Engl, Hanke, Kunisch 1989 Inverse Problems

- $f := 16$, $u(x) = 8x(1 - x)$
- $u_n = u + e_n$, with e_n , $n \geq 2$, given by

$$e_n = \begin{cases} -n^{-3/4}(2x)^{2n}, & x \leq 1/2, \\ -n^{-3/4}(2 - 2x)^{2n}, & x > 1/2 \end{cases}$$

- The unique sol. in $D(F)$ is $q^\dagger \equiv 0$ and $q_n = e_n''/(u + e_n)$
- $u_n \rightarrow u$ in $L^2(\Omega)$. but $\|q_n\|_{L^2(\Omega)} \rightarrow \infty$, i.e., $q_n \not\rightarrow q^\dagger$ in L^2

Thus q does not depend continuously on $F(q)$ as a mapping from $L^2(\Omega)$ to $L^2(\Omega)$



What about Tikhonov approach ?

$$J_{\alpha}(q) = \|F(q) - g\|^2 + \alpha \|q\|^2$$

with $F(q) : q \rightarrow u(q)$, with q belonging to $\mathcal{A}$

$$\mathcal{A} = \{q \in L^{\infty}(\Omega) : q \geq 0, q \leq c\}.$$

What about the existence of a solution to the problem ?



argument:

- $J_\alpha(q) \geq 0 \Rightarrow \exists$ a minimizing sequence $\{q^n\} \subset \mathcal{A}$

$$\lim_{n \rightarrow \infty} J_\alpha(q^n) = \inf_{q \in \mathcal{A}} J_\alpha(q)$$

- The sequence $\{q^n\}$ is uniformly bdd. in L^2
 $\Rightarrow \exists$ a subsequence $\{q^{n_k}\}$ converging weakly in $L^2(\Omega)$
- $F(q^{n_k}) \rightarrow F(q^*)$ weakly in $L^2(\Omega)$?
- weak lower semi-continuity of norm $\Rightarrow \exists$ a minimizer

weak continuity ?



weak continuity

The forward operator $F(q)$ is weakly continuous from $L^2(\Omega)$ to $L^2(\Omega)$.

Let $q^n \rightarrow q^*$ weakly in $L^2(\Omega)$

$$\int_{\Omega} \nabla u^n \cdot \nabla v dx + \int_{\Omega} q^n u^n v dx = \int_{\Omega} f v dx \quad \forall v \in H_0^1(\Omega)$$

Taking $v = u^n \Rightarrow \|u^n\|_{H^1(\Omega)} \leq c$

$\Rightarrow \exists$ a subsequence $\{u^n\}$ converges weakly to u^* in $H^1(\Omega)$

$$\begin{aligned} \int_{\Omega} \nabla u^n \cdot \nabla v dx &\rightarrow \int_{\Omega} \nabla u^* \cdot \nabla v dx \quad \forall v \in H_0^1(\Omega) \\ \underbrace{\int_{\Omega} q^n u^n v dx}_{\text{prod. of weakly conv. seq.}} &= \int_{\Omega} q^n u^* v dx + \int_{\Omega} q^n (u^n - u^*) v dx \end{aligned}$$

prod. of weakly conv. seq.



$u^*, v \in H_0^1(\Omega) \Rightarrow u^*, v \in L^4(\Omega)$ (Sobolev embedding) for $d = 2, 3$
and the embedding into $L^p(\Omega)$, $p < 4$ is compact

■ $u^*, v \in L^4(\Omega) \Rightarrow u^* v \in L^2(\Omega) + \text{weak conv. of } q^n \text{ in } L^2(\Omega)$

$$\int_{\Omega} q^n u^* v dx \rightarrow \int_{\Omega} q^* u^* v dx$$

■ $u^n \rightarrow u^*$ in $L^2(\Omega) + \text{bbd of } q^n$

$$\left| \int_{\Omega} q^n (u^n - u^*) v dx \right| \leq \|q^n\|_{L^\infty(\Omega)} \|u^n - u^*\|_{L^2(\Omega)} \|v\|_{L^2(\Omega)} \rightarrow 0$$

■ the limit u^* satisfies ($u^* = u(q^*)$)

$$\int_{\Omega} \nabla u^* \cdot \nabla v dx + \int_{\Omega} q^* u^* v dx = \int_{\Omega} f v dx \quad \forall v \in H_0^1(\Omega)$$

■ Every conv. subseq. of $\{u^n\}$ has a sub. seq. converging to $u(q^*)$
 $\Rightarrow$ the whole sequence converges weakly to $u(q^*)$ in $H^1(\Omega)$