

STRUCTURING CAUSAL TREE MODELS WITH CONTINUOUS VARIABLES *

Lei Xu

Department of Automation
Tsinghua University
Beijing China

Judea Pearl

Cognitive Systems Laboratory
Computer Science Department
UCLA, Los Angeles, CA. 90024-1600

ABSTRACT

This paper considers the problem of invoking auxiliary, unobservable variables to facilitate the structuring of causal tree models for a given set of continuous variables. Paralleling the treatment of bi-valued variables in [Pearl 1986], we show that if a collection of coupled variables are governed by a joint normal distribution and a tree-structured representation exists, then both the topology and all internal relationships of the tree can be uncovered by observing pairwise dependencies among the observed variables (i.e., the leaves of the tree). Furthermore, the conditions for normally distributed variables are less restrictive than those governing bi-valued variables. The result extends the applications of causal tree models which were found useful in evidential reasoning tasks.

I. INTRODUCTION

Belief networks are directed acyclic graphs in which nodes represent random variables, the arcs signify direct dependencies between the linked variables, and the strengths of these dependencies are quantified by conditional probabilities. Belief networks can be used to represent the generic knowledge of a domain, as well as inference engines that manipulate this knowledge in evidential reasoning applications [Pearl 1986, 1988]. In particular, it was shown that in a singly-connected (e.g. tree-structured) network, beliefs can be updated coherently by local propagation through a network of parallel and autonomous processors, and that equilibrium is guaranteed to be reached in time proportion to the network diameter.

Since the efficacy of the scheme depends on the singly-connected structure of the network, Pearl [1986] has proposed a preprocessing approach which introduces auxiliary variables and permanently turns multiply-connected belief networks into a tree. In [Pearl, 1988], it is further argued that the introduction of auxiliary variables into causal models constitutes an important component of human learning, aimed at attaining the features of modularity and

* This work was supported in part by the National Science Foundation Grant, DCR 83-13875.

autonomy that are unique to tree representations. Naturally, the question arises what class of multiply-connected networks can be reconfigured as trees. In [Pearl, 1986] it is shown that if all variables are bi-valued and if there exists a decomposition into a tree-structured network with auxiliary variables, then the topology of the tree can be uncovered uniquely from the observed correlations between pairs of variables. However, if the variables are not bi-valued, the problem is more complicated and remains an open question.

In this paper, we extend the method of structuring causal trees from bi-valued to continuous normal variables. We show that if all the variables are normally distributed and if the activities of the visible variables are governed by a tree-decomposable joint normal distribution, then the tree can be structured from the observed correlations between pairs of variables. Moreover, the conditions for normally distributed variables to be tree-decomposable are less restrictive than the corresponding conditions for bi-valued variables.

II. THEORETICAL BACKGROUND

A. Nomenclature and Problem Statement

Let X_1, X_2, \dots, X_n be random variables from a n -dimensional joint normal distribution:

$$f(x_1, x_2, \dots, x_n) = (2\pi)^{-\frac{n}{2}} (\det \Sigma_n)^{-\frac{1}{2}} \exp \left[-\frac{1}{2} (x_n - \mu_n)^t \Sigma_n^{-1} (x_n - \mu_n) \right], \quad (1)$$

where $x_n = (x_1, x_2, \dots, x_n)^t$, $\mu_n = E x_n$ is the mean vector and $\Sigma_n = E(x_n - \mu_n)(x_n - \mu_n)^t$ is the covariance matrix of x_n .

Analogous with Section 3.2 in [Pearl, 1986], we can ask if $f(x_1, x_2, \dots, x_n)$ can be represented as a marginal of an $n+1$ dimensional normal distribution of $x_{n+1} = (w, x_n^t)^t$ such that the X_i 's are conditionally independent given W , i.e.

$$f(x_1, x_2, \dots, x_n) = \int_{-\infty}^{+\infty} f_s(x_1, x_2, \dots, x_n, w) dw \quad (2)$$

$$f_s(x_1, x_2, \dots, x_n, w) = \prod_{i=1}^n f_s(x_i | w) f(w), \quad (3)$$

where $f_s(x_i | w)$, $i = 1, \dots, n$ relate each X_i to the central hidden variable W (see Figure 1).




Figure 1

If the decomposition in (2) is possible, we name f_s a *star-distribution* and call f *star-decomposable*.

Instead of one hidden variable, W , we can use m hidden variables ($m \leq n - 2$) to form a tree-like structure (see Figure 2), in which each triplet of leaves forms a star, but the central variable may differ from triplet to triplet.




Figure 2

A normal distribution $f(x_1, x_2, \dots, x_n)$ is said to be *tree-decomposable* if it is a marginal of an $n + m$ dimensional normal distribution:

$$f_t(x_1, x_2, \dots, x_n, w_1, w_2, \dots, w_m) \quad m \leq n - 2 \quad (4)$$

that maps into a tree; i.e., W_1, W_2, \dots, W_m correspond to the internal nodes of a tree and X_1, X_2, \dots, X_n to its leaves.

In this paper we assume that a) each W has at least three neighbors and b) $f(x|y) \neq f(x)$ for each $X, Y \in \{X_1, X_2, \dots, X_n, W_1, W_2, \dots, W_m\}$, i.e., there are genuine dependencies between the linked variables (otherwise the tree can be decomposed into a forest).

The problem is whether $f_t(x_1, x_2, \dots, x_n, w_1, w_2, \dots, w_m)$ can be recovered given a tree-decomposable normal distribution $f(x_1, x_2, \dots, x_n)$.

B. A Theorem on Normal Distributions

It is well known (e.g., [Gigi 1977]) that the covariance matrix Σ of a normal random vector $\mathbf{x} = (x_1, x_2, \dots, x_p)^t$ is a $p \times p$ diagonal matrix, iff the components of \mathbf{x} represent independent normal random variables. Moreover,

Theorem 1: Let $\mathbf{x} = (\mathbf{x}_{(1)}^t, \mathbf{x}_{(2)}^t)^t$, $\mathbf{x}_{(1)} = (x_1 \cdots x_q)$, $\mathbf{x}_{(2)} = (x_{q+1} \cdots x_p)$.

Let $\mu = E\mathbf{x}$ be similarly partitioned as $\mu = (\mu_{(1)}, \mu_{(2)})^t$ and let Σ be partitioned as

$$\Sigma = \begin{Bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{Bmatrix} \quad (5)$$

where Σ_{11} is the $q \times q$ upper left-hand corner submatrix of Σ . If \mathbf{x} is normally distributed with mean μ and covariance matrix Σ , then

- The vectors $\mathbf{x}_{(1)}$ and $\mathbf{x}_{(2)} - \Sigma_{21} \Sigma_{11}^{-1} \mathbf{x}_{(1)}$ are independently normally distributed with means $\mu_{(1)}$, $\mu_{(2)} - \Sigma_{21} \Sigma_{11}^{-1} \mu_{(1)}$, and covariance matrices Σ_{11} , $\Sigma_{22} - \Sigma_{21} \Sigma_{11}^{-1} \Sigma_{12}$, respectively.
- The marginal distribution of $\mathbf{x}_{(1)}$ is q -variate normal with mean $\mu_{(1)}$ and covariance matrix Σ_{11} .
- The conditional distribution of $\mathbf{x}_{(2)}$ given $\mathbf{x}_{(1)}$ is normal with mean $\mu_{(2)} + \Sigma_{21} \Sigma_{11}^{-1} (\mathbf{x}_{(1)} - \mu_{(1)})$ and covariance matrix $\Sigma_{22} - \Sigma_{21} \Sigma_{11}^{-1} \Sigma_{12}$.

The proof of the theorem is given in [Gigi, 1977, pp. 51-53] and it will be used in the next section.

III. STAR-DECOMPOSABLE TRIPLETS OF GAUSSIAN VARIABLES

Let $f(x_1, x_2, x_3)$ be a 3-dimensional joint normal distribution as in (1), with mean $\mu = (\mu_1 \mu_2 \mu_3)^t$ and covariance matrix

$$\Sigma = \begin{vmatrix} \sigma_{11} & \sigma_{12} & \sigma_{13} \\ \sigma_{21} & \sigma_{22} & \sigma_{23} \\ \sigma_{31} & \sigma_{32} & \sigma_{33} \end{vmatrix} \quad \text{and} \quad \sigma_{ij} = \sigma_{ji}, \quad i, j = 1, 2, 3. \quad (6)$$

If $f(x_1, x_2, x_3)$ is star-decomposable, then it is a marginal of a 4-dimensional joint normal distribution $f_s(w, x_1, x_2, x_3)$ with mean $\mu_s = (\mu_w, \mu')^t$ and covariance matrix

$$\Sigma_s = \begin{array}{c|ccc} & \sigma_{ww} & \sigma_{1w} & \sigma_{2w} & \sigma_{3w} \\ \hline \sigma_{w1} & & & & \\ \sigma_{w2} & & & \Sigma & \\ \sigma_{w3} & & & & \end{array}$$

and

$$\sigma_{wi} = \sigma_{iw}, \quad i = 1, 2, 3$$

$$f(w, x_1, x_2, x_3) = f(x_1, x_2, x_3 | w) f(w) \quad (7)$$

$$f(x_1, x_2, x_3 | w) = f(x_1 | w) f(x_2 | w) f(x_3 | w). \quad (8)$$

Theorem 1 states that $f(x_1, x_2, x_3 | w)$, $f(w)$ and $f(x_i | w)$'s are also normal distributions, and the mean vector and covariance matrix of $f(x_1, x_2, x_3 | w)$ are given by

$$\mu_{1:2:3|w} = \mu - \sigma_{ww}^{-1} \begin{bmatrix} \sigma_{w1} & \sigma_{w2} & \sigma_{w3} \end{bmatrix}^t (w - \mu_w) \quad (9)$$

$$\Sigma_{1:2:3|w} = \Sigma - \sigma_{ww}^{-1} \begin{bmatrix} \sigma_{w1} & \sigma_{w2} & \sigma_{w3} \end{bmatrix}^t \begin{bmatrix} \sigma_{w1} & \sigma_{w2} & \sigma_{w3} \end{bmatrix}. \quad (10)$$

Additionally, the conditional independence stated in (8) implies that $\Sigma_{1:2:3|w}$ must be a diagonal matrix, thus

$$\sigma_{ij} - \sigma_{wi} \sigma_{jw} / \sigma_{ww} = 0, \quad i \neq j \text{ and } i, j = 1, 2, 3. \quad (11)$$

and

$$\sigma_{ii} - \sigma_{iw}^2 / \sigma_{ww} > 0, \quad i = 1, 2, 3. \quad (12)$$

Using the correlation coefficients defined as

$$\rho_{ij} = \sigma_{ij} / (\sigma_{ii} \sigma_{jj})^{1/2} \quad (13)$$

(11) and (12) can be written as

$$\rho_{ij} = \rho_{iw} \rho_{jw}, \quad \text{for all } i, j \quad (14)$$

$$\rho_{iw}^2 \leq 1 \quad \text{for all } i. \quad (15)$$

Solving (14) for ρ_{iw} , we obtain

$$\rho_{1w} = (\rho_{12}\rho_{13} / \rho_{23})^{1/2} \quad \rho_{2w} = (\rho_{12}\rho_{23} / \rho_{13})^{1/2} \quad \rho_{3w} = (\rho_{13}\rho_{23} / \rho_{12})^{1/2} \quad (16)$$

The requirement that the ρ_{iw} 's must be real numbers with magnitude not exceeding 1, yields the following two conditions for $f(x_1, x_2, x_3)$ to be star-decomposable:

- a. $\rho_{12}, \rho_{13}, \rho_{23}$ are all positive, or two are negative and one is positive. In other words, the triplet (x_1, x_2, x_3) is positively correlated.
- b. $\rho_{jk} \geq \rho_{ji}\rho_{ik}$ for all $i, j, k \in \{1, 2, 3\}$ and $i \neq j \neq k$

Summarizing the analysis above, we obtain Theorem 2.

Theorem 2:

1. A necessary and sufficient condition for three random variables with a joint normal distribution to be star-decomposable is that the correlation coefficients satisfy the inequalities:

$$\rho_{jk} \geq \rho_{ji}\rho_{ik} \quad \rho_{ij} \rho_{ji} \rho_{ki} \geq 0 \quad (17)$$

for all $i, j, k \in \{1, 2, 3\}$, and $i \neq j \neq k$.

2. $f(x_i | w) \sim N(\mu_i | w, \sigma_i | w)$, $i = 1, 2, 3$ are specified by the parameters $\sigma_i | w = \sigma_{ii}(1 - \rho_{iw}^2) = \sigma_{ii}(1 - \rho_{ji}\rho_{ik} / \rho_{jk})$
 $\mu_i | w = \mu_i - \sigma_{wi}(w - \mu_w) / \sigma_{ww} = \mu_i - \rho_{wi} \sqrt{\frac{\sigma_{ij}}{\sigma_{ww}}} (w - \mu_w)$ and
 $f(w) \sim N(\mu_w, \sigma_w)$, where $\sigma_{ww} > 0$ and μ_w may be chosen arbitrarily.

Part 2 of Theorem 2 can be proved from (9), (10) and (11) combined with (13) and (16).

In a manner similar to [Pearl, 1986] we may pose the following problem. Suppose $f(x_1, x_2, x_3)$ is an arbitrary distribution (not necessarily normal), can it be approximated by a star-decomposable normal distribution $f_s(x_1, x_2, x_3)$ which will have the same covariance matrix as f . The answer is implied by Theorem 2, and stated in Theorem 3.

Theorem 3: A necessary and sufficient condition for the second order dependencies among the triplet X_1, X_2, X_3 to support a star-decomposable normal joint distribution is that all the correlation coefficients obey the triangle inequality:

$$\rho_{12}\rho_{13}\rho_{23} \geq 0 \quad \text{and} \quad \rho_{jk} \geq \rho_{ji}\rho_{ik} \quad \text{for all } i, j, k \in \{1, 2, 3\}, i \neq j \neq k.$$

Discussion:

1. Comparing Theorem 2 and 3 to those of [Pearl, 1986], we see that the conditions for a normally distributed triplet X_1, X_2, X_3 to be star-decomposable are less restrictive than those for bi-valued variables; there is no restriction corresponding to the 3rd order constraint imposed on P_{ijk} in [Pearl, 1986].

2. Part 2 of Theorem 2 illustrates that if the conditions (17) are satisfied, the densities $f(w)$, $f(x_i | w)$'s can be obtained in closed form, depending on the selection of μ_w, σ_w . In the case of star-decomposable triplets, one may conveniently set $\sigma_w^2 = 1$ and $\mu_w = 0$ i.e., let the inner node have a standard normal distribution. For tree-decomposable structures with more than three variables, the selection of μ_w, σ_w for the intermediate variables should be made in a consistent manner, as shown in the next section.
3. To ensure that $n (>3)$ variables of a joint normal distribution be star-decomposable, (14) and (15) must lead to a consistent solution for all the ρ_{iw} 's.
4. A simple (degenerate) example of triplets which always satisfy (17) is given by 1st-order Markov variables X_1, X_2, X_3 , governed by the covariance matrix

$$\Sigma = \begin{vmatrix} \sigma_{11} & \rho\sqrt{\sigma_{11}\sigma_{22}} & \rho^2\sqrt{\sigma_{11}\sigma_{33}} \\ \rho\sqrt{\sigma_{11}\sigma_{22}} & \sigma_{22} & \rho\sigma_{22}\sigma_{33} \\ \rho^2\sqrt{\sigma_{11}\sigma_{33}} & \rho\sqrt{\sigma_{22}\sigma_{33}} & \sigma_{33} \end{vmatrix} \quad 1 > \rho > 0$$

Since $\rho_{12} = \rho$, $\rho_{13} = \rho^2$ and $\rho_{32} = \rho$,

we have

$$\begin{aligned} \rho_{12} &> \rho_{13} \rho_{32} \\ \rho_{23} &> \rho_{12} \rho_{13} \\ \rho_{13} &= \rho_{12} \rho_{23} \end{aligned}$$

In this case the central variable W coincides with X_2 .

IV. STRUCTURING CAUSAL TREES

Consider the 4 possible topologies of 4-tuple of leaves in a tree, as given in Figure 3.




Figure 3

The topologies differ in the identity of the triplets which share a common center, e.g., in the topology of Figure 3 (a), [(1, 2, 3), (1, 2, 4)] share a common center; so does the pair [(1, 3, 4), (2, 3, 4)]. It follows that the star-decomposition of either triplet (1, 2, 3) or (1, 2, 4) should yield the same values of σ_w, μ_w . This means that the correlation coefficients of the 4 variables with a topology of Figure 3 (a) should satisfy (16),

$$\rho_{1w}^2 = \rho_{12}\rho_{13} / \rho_{23} \quad \text{for } (1, 2, 3)$$

$$\rho_{1w}^2 = \rho_{12}\rho_{14} / \rho_{24} \quad \text{for } (1, 2, 4)$$

Thus,

$$\rho_{12}\rho_{13} / \rho_{23} = \rho_{12}\rho_{14} / \rho_{24}$$

or

$$\rho_{13}\rho_{24} = \rho_{14}\rho_{23} \quad (18)$$

(Enforcing the equality of ρ_{2w}^2 for (1, 2, 3) and (1, 2, 4), would yield the same equation). Similarly, from the pair [(1, 3, 4), (2, 3, 4)], we also obtain (18) (note $\rho_{ij} = \rho_{ji}$). Thus, the equality in (18) may be taken as the essential condition identifying the topology of Figure 3(a).

The equalities characterizing the 4 possible topologies of Figure 3 are given in table 1.

Fig. 3(a)	Fig. 3(b)	Fig. 3(c)	Fig. 3(d)
$\rho_{13}\rho_{24} = \rho_{14}\rho_{23}$	$\rho_{12}\rho_{34} = \rho_{14}\rho_{23}$	$\rho_{12}\rho_{34} = \rho_{13}\rho_{24}$	two equalities holding simultaneously

Table 1. The Characteristic Equalities of the 4 Topologies in Figure 3

Table 1 may be taken as a tool for both testing if a 4-tuple is tree-decomposable and, if so, deciding its topology. Since the basic test to decide the topology of any 4-tuple is the same as that of binary variables, the topology of the entire tree of n leaves can be decided by the procedure described in [Pearl and Tarsi, 1986].

The table yields the following condition for tree-decomposability:

Theorem 4: A necessary condition for a joint normal distribution to be tree-decomposable is that, for every set of four variables $\{X_i, X_j, X_k, X_l\}$, the equality

$$\rho_{\alpha\beta}\rho_{\gamma\sigma} = \rho_{\alpha\gamma}\rho_{\beta\sigma}$$

be satisfied for at least one permutation $(\alpha, \beta, \gamma, \sigma)$ of (i, j, k, l) .

The next stage is to determine all the $f(x_i|w)$ and $f(w_j|w_k)$ functions. The functions $f(x_i|w_j)$ assigned to the peripheral branches of the tree are determined directly from the star-decomposition of triplets involving adjacent leaves, and these can be used to determine the functions $f(w_j|w_k)$ connecting the hidden variables. In Figure 4, for example, the star-decomposition of $f(x_1, x_2, x_3)$ yields $f(x_1|w_1)$, $f(x_2|w_1)$, while $f(x_1|w_2)$ can be obtained from the star-decomposition of (X_1, X_3, X_6) . These are sufficient for determining $f(w_1|w_2)$, via

$$f(x_1|w_2) = \int_{w_1} f(x_1|w_1) f(w_1|w_2) dw_1 \quad (19)$$




Figure 4

Let the mean and variance of $f(x_1|w_2)$, $f(x_1|w_1)$ and $f(w_1|w_2)$ be denoted by $\mu_{x_1|w_2}$, $\sigma_{x_1|w_2}^2$, $\mu_{x_1|w_1}$, $\sigma_{x_1|w_1}^2$ and $\mu_{w_1|w_2}$, $\sigma_{w_1|w_2}^2$ respectively, and let the mean vector and covariance matrix of (X_1, W_1, W_2) be $[\mu_{x_1}, \mu_{w_1}, \mu_{w_2}]$ and

$$\begin{vmatrix} \sigma_{x_1}^2 & \sigma_{x_1 w_1} & \sigma_{x_1 w_2} \\ \sigma_{w_1 x_1} & \sigma_{w_1}^2 & \sigma_{w_1 w_2} \\ \sigma_{w_2 x_1} & \sigma_{w_1 w_2} & \sigma_{w_2}^2 \end{vmatrix}$$

We will show that $\sigma_{w_1|w_2}$, $\mu_{w_1|w_2}$ (thus $f(w_1|w_2)$) may be determined from $f(x_1|w_2)$ and $f(x_1|w_1)$. From Theorem 1, the mean vector and covariance matrix of $f(x_1|w_1) f(w_1|w_2) = f(x_1, w_1|w_2)$ and $f(x_1, w_2|w_1)$ are given by

$$\Sigma_{x_1 w_1 | w_2} = \begin{Bmatrix} \sigma_{x_1}^2 & \sigma_{x_1 w_1} \\ \sigma_{w_1 x_1} & \sigma_{w_1}^2 \end{Bmatrix} - \frac{1}{\sigma_{w_2}^2} \begin{Bmatrix} \sigma_{w_2 x_1} \\ \sigma_{w_2 w_1} \end{Bmatrix} \begin{Bmatrix} \sigma_{w_2 x_1} & \sigma_{w_2 w_1} \end{Bmatrix}, \quad (20)$$

$$\Sigma_{x_1 w_2 | w_1} = \begin{Bmatrix} \sigma_{x_1}^2 & \sigma_{w_2 x_1} \\ \sigma_{x_1 w_2} & \sigma_{w_2}^2 \end{Bmatrix} - \frac{1}{\sigma_{w_1}^2} \begin{Bmatrix} \sigma_{w_1 x_1} \\ \sigma_{w_1 w_2} \end{Bmatrix} \begin{Bmatrix} \sigma_{w_1 x_1} & \sigma_{w_1 w_2} \end{Bmatrix},$$

$$\mu_{x_1 w_1 | w_2} = \begin{Bmatrix} \mu_{x_1} \\ \mu_{w_1} \end{Bmatrix} - \frac{1}{\sigma_{w_2}^2} \begin{Bmatrix} \sigma_{w_2 x_1} \\ \sigma_{w_2 w_1} \end{Bmatrix} (w_2 - \mu_{w_2}), \quad (21)$$

$$\mu_{x_1 w_2 | w_1} = \begin{Bmatrix} \mu_{x_1} \\ \mu_{w_2} \end{Bmatrix} - \frac{1}{\sigma_{w_1}^2} \begin{Bmatrix} \sigma_{w_1 x_1} \\ \sigma_{w_1 w_2} \end{Bmatrix} (w_1 - \mu_{w_1}).$$

Therefore,

$$\sigma_{x_1 | w_2}^2 = \sigma_{x_1}^2 - \sigma_{w_2 x_1}^2 / \sigma_{w_2}^2 \quad (22)$$

$$\sigma_{w_1 | w_2}^2 = \sigma_{w_1}^2 - \sigma_{w_2 w_1}^2 / \sigma_{w_2}^2$$

$$\sigma_{x_1 | w_1}^2 = \sigma_{x_1}^2 - \sigma_{w_1 x_1}^2 / \sigma_{w_1}^2$$

$$\mu_{x_1 | w_2} = \mu_{x_1} - \frac{\sigma_{w_2 x_1}}{\sigma_{w_2}^2} (w_2 - \mu_{w_2}) \quad (23)$$

$$\mu_{w_1 | w_2} = \mu_{w_1} - \frac{\sigma_{w_2 w_1}}{\sigma_{w_2}^2} (w_2 - \mu_{w_2})$$

$$\mu_{x_1 | w_1} = \mu_{x_1} - \frac{\sigma_{w_1 x_1}}{\sigma_{w_1}^2} (w_1 - \mu_{w_1}).$$

It is to be noted that $f(x_1 w_2 | w_1) = f(x_1 | w_1) f(w_2 | w_1)$, hence, $\Sigma_{x_1 w_2 | w_1}$ is a diagonal matrix which leads to

$$\sigma_{w_2 x_1} = \sigma_{w_1 x_1} \sigma_{w_2 w_1} / \sigma_{w_1}^2 \quad (24)$$

As mentioned in Theorem 2, the mean and variance of the first inner variable may be assigned arbitrarily. We set $\sigma_{w_1}^2 = 1$ and $\mu_{w_1} = 0$, then join (22) and (24) and keep in mind that $\sigma_{x_1 | w_1}^2, \sigma_{x_1}^2, \sigma_{x_1 | w_2}^2$ and $\mu_{x_1 | w_2}, \mu_{x_1}, \mu_{x_2 | w_1}$ are known from $f(x_1 | w_1), f(x_1 | w_2)$ and $f(x_1)$, and $\sigma_{w_1 | w_2}^2, \sigma_{w_2}^2, \sigma_{w_1 w_2}, \sigma_{w_1 x_1}$ may be solved. In turn,

$\mu_{w_2}, \mu_{w_1|w_2}$ can be solved by (23). Consequently, $f(w_1|w_2)$ is determined by $\sigma_{w_1|w_2}, \mu_{w_1|w_2}$; $f(w_2)$ is determined by $\sigma_{w_2}^2, \mu_{w_2}$. In the same manner, $f(w_3|w_2)$, and $f(w_4|w_2)$ can also be determined.

V. CONCLUSIONS

The paper extends the auxiliary-variable method of constructing causal tree representations to continuous variables. It shows that if the visible variables are governed by a tree-decomposable joint normal distribution, then the tree can be structured from the observed correlations between pairs of variables. Furthermore, the conditions for tree-decomposable normal distribution are less restrictive than those of bi-valued variables. The results should extend the applicability of causal tree models to evidential reasoning tasks involving continuous signals and can also be used to discover the underlying causal structure behind ill understood phenomena.

REFERENCES

- [Gigi, 1977] Gigi, N. C., *Multivariate Statistical Inference*, Academic Press, New York, 1977.
- [Pearl, 1986] Pearl, J., Fusion, propagation and structuring in belief networks, *Artificial Intelligence*, Vol. 29, No. 3, September 1986, pp. 241-288.
- [Pearl, 1988] Pearl, J., *Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference*, Morgan Kaufmann, San Mateo, CA.
- [Pearl and Tarsi, 1986] Pearl, J., and Tarsi, M., Structuring causal trees, *Journal of Complexity*, 2 (no. 1), pp. 60-77.